







VITAMIN K AMPOULE (ORAL)

Konakion® MM Paediatric Injection 2 mg/0.2 mL

Patient Information Leaflet

This leaflet illustrates the proper method of breaking and administering Vitamin K orally from the ampoule.

The glass ampoules are constructed to ensure a clean break provided the following steps are followed:

<p>Step 1:</p> <ul style="list-style-type: none">• Shake the ampoule until all the solution is at the bottom.• The solution should be clear; do not use if cloudy. 	<p>Step 2:</p> <ul style="list-style-type: none">• Hold the bottom part of the ampoule between the thumb and first finger of one hand.• Ensure that the spot on the ampoule is facing towards you. 
<p>Step 3:</p> <ul style="list-style-type: none">• If possible, cover the top of the ampoule with a tissue or towel before holding the top between the thumb and first finger of your other hand. 	<p>Step 4:</p> <ul style="list-style-type: none">• Snap the top of the ampoule off by pushing away from the side with the spot. 
<p>Step 5:</p> <ul style="list-style-type: none">• Place the dispenser (provided together with the ampoule) vertically into the ampoule.• Withdraw the solution into the dispenser until it reaches the correct marking on the dispenser.• Hold the dispenser against the light to check whether the solution is clear and also ensure that no glass pieces are stuck to the tip of the dispenser.• If unsure, expel the solution into a small cup to check for any glass pieces – if the solution is clear, withdraw again from the cup into the dispenser. 	<p>Step 6:</p> <ul style="list-style-type: none">• Administer the contents of the dispenser directly into the mouth. 

Storage:

Store in a cool, dry place (room temperature), away from light.