

Contact Information

NNI@TTSH

Tan Tock Seng Hospital, NNI Block, Neuroscience Clinic
11 Jalan Tan Tock Seng, Singapore 308433
Main Tel: (65) 6357 7153
Appt. Tel: (65) 6330 6363
Email: appointments@nni.com.sg
Website: www.nni.com.sg



NNI@SGH

Singapore General Hospital, Block 3, Clinic L
Outram Road, Singapore 169608
Main Tel: (65) 6222 3322
Appt. Tel: (65) 6321 4377
Email: appointments@sgh.com.sg
Website: www.nni.com.sg



NNI@CGH

Changi General Hospital
2 Simei Street 3
Singapore 529889
Appt. Tel: (65) 6850 3333

NNI@KKH

KK Women's and
Children's Hospital
100 Bukit Timah Road
Singapore 229899
Appt. Tel: (65) 6294 4050

NNI@KTPH

Khoo Teck Puat Hospital
90 Yishun Central
Singapore 768828
Appt. Tel: (65) 6555 8828


NNI@SKH

Sengkang General Hospital
110 Sengkang East Way
Singapore 544886
Appt. Tel: (65) 6930 6000




Scan the QR code to learn
more about other
Neuroscience conditions

Neurosurgery Department



The National Neuroscience Institute operates out of
two main campuses (TTSH, SGH) and
four partner hospitals (CGH, KKH, KTPH, SKH).



Brain Tumours

Acoustic Neuroma

Brochure content serves as a guide only
Seek the advice of your doctor for more details

Information correct as of April 2020



National
Neuroscience Institute
SingHealth

Understanding Acoustic Neuromas

An acoustic neuroma is a non-cancerous growth that occurs near the facial nerve. They are slow-growing and may remain the same size for years for some patients. In a small number of patients, the tumour becomes smaller with time.

Signs of Acoustic Neuromas

Signs vary and include:

- One-sided deafness, with noise in the affected ear (tinnitus). The deafness may be gradual or sudden and is experienced by 90% of patients
- Inability to understand speech i.e. one can hear sounds but cannot understand what is being said
- Unsteady gait and poor balance
- Facial numbness and weakness
- Swallowing problems

Diagnosing Acoustic Neuromas

Auditory tests can reveal loss of hearing and the inability to understand speech.

A Computed Tomography (CT) or Magnetic Resonance Imaging (MRI) scan can show the presence of an acoustic neuroma, even those that are still in the internal ear canal (Figure 1).



Figure 1
Area inside the inner ear where tumours grow

Risk Factors

Those with rare genetic defects e.g. tumours growing on nerve tissues (neurofibromatosis) may be at a higher risk.

Treating Acoustic Neuromas

Surgery

Surgery may be required for patients with very large tumours causing severe brain compression and increased brain pressure.

A neurosurgeon and sometimes a ear, nose and throat surgeon will determine if surgery is suitable, depending on the size and location of the tumour, and the health of the patient.

During surgery, fat or muscle may be taken from the abdomen or thigh to close the wound. After surgery, the patient will usually spend one to several days in the intensive care unit for monitoring and treatment.

Depending on the location of the tumour, side effects of the surgery may include hearing loss, facial weakness, paralysis, double vision, swallowing problems, mouth dryness, and unsteadiness.

The likelihood of unexpected complications is generally low but include infection, bleeding, stroke, seizures, paralysis of limbs, coma and death.

Radiation Therapy

Smaller tumours are usually treated by radiation. A high and precise dose of radiation is aimed at the tumour, with no or low damage to surrounding brain structures.

Support for Brain Tumours

Brain Tumour Society Singapore (BTSS)

BTSS is a community of brain tumour patients, caregivers and survivors. The BTSS provides community support and resources such as befrienders, financial assistance and public education. Started by brain cancer survivors, BTSS meets once a month so that members can share experiences and advice on how to cope with the disease.

For more information, visit www.braintumoursociety.org.sg.